



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

233. (June, 1915.) Proposed by V. M. SPUNAR, Chicago, Illinois.

Solve in rational numbers $x^2 + y^2 = a^2$, $xy = m/n$, when m and n are integers and relatively prime to each other.

SOLUTION BY A. H. HOLMES, Brunswick, Maine.

Putting $x = (p^2 - q^2)/(2p \pm 2q + 1)$, $y = (2pq)/(2p \pm 2q + 1)$, we shall have

$$a = (p^2 + q^2)/(2p \pm 2q + 1), \quad m = 2pq(p^2 - q^2),$$

and $n = (2p \pm 2q + 1)^2$ except in the cases where $p = q + 1$, $2q + 1$, or $5q + 1$, in which cases $n = 2p \pm 2q + 1$.

For smallest values put $q = 1$, $p = 2$, using the minus sign of the denominator. Then $x = 1$, $y = 4/3$, $a = 5/3$, $m = 4$, and $n = 3$.

Also solved by H. N. CARLETON.

QUESTIONS AND DISCUSSIONS.

SEND ALL COMMUNICATIONS TO U. G. MITCHELL, University of Kansas, Lawrence.

DISCUSSIONS.

I. RELATING TO AN INSTRUCTIVE PROBLEM IN ATTRACTION.

BY EDWIN BIDWELL WILSON, Massachusetts Institute of Technology.

In B. O. Pierce's *Newtonian Potential Function* (3d ed.) there is found on page 28 this exercise:

Show that the attraction at the focus of a segment of a paraboloid of revolution bounded by a plane perpendicular to the axis at a distance b from the vertex is of the form $4\pi\rho a \log(1 + b/a)$.

This problem is also found in other texts. The answer is very easy to get; it is also wrong. One way in which the answer may be found is this:

The attraction of a disc of mass M and radius r at a point on its axis at a distance c from the center is

$$\frac{2M}{r^2} \left(1 - \frac{c}{\sqrt{c^2 + r^2}} \right).$$

If the equation of the revolved parabola is $4ay = x^2$, the attraction of a segment at the focus is

$$\frac{2\pi\rho \cdot 4ay}{4ay} \left(1 - \frac{y - a}{\sqrt{(y - a)^2 + 4ay}} \right) dy,$$

and the total attraction is

$$A = 2\pi\rho \int_0^b \left(1 - \frac{y - a}{\sqrt{(y - a)^2 + 4ay}} \right) dy.$$

To the careless person this gives

$$A = 2\pi\rho \int_0^b \frac{2ady}{y + a} = 4\pi\rho a \log(1 + b/a)$$